Scalar-mediated graviton mass generation in spontaneous scalarized DNS binaries

Jing Wang*
School of Physics and Astronomy, Sun Yat-Sen University, Guangzhou, 510275, P. R. China

We propose the mechanism that gravitons radiated from double neutron star (DNS) binaries acquire a mass term, owing to a mediation of massive scalar field. A scalarized DNS system, whose dynamics is governed by scalar-tensor theory, undergoes a spontaneous symmetry breaking, arising from the coupling of scalarized NSs to the gravitational scalar field and the scalar fluctuations. The resulted Higgs-like potential endows the scalar field with a mass, where the scalar fluctuation plays a role of the Higgs field. The Yukawa-like correction to the Newtonian potential of the binary subsequently induces a mass term for gravitons during their propagations. We estimate the mass of gravitons for 8 detected DNS binaries with more precise mass measurements, which is several times of $10^{-24}ev/c^2$ and mildly varies according to the compactness of scalarized NSs and the separation between them. Because the gravitational massive scalar mode is suppressed on a scale of 10^9 m by the Yukawa-like potential, we claim that the massive gravitons doesn't suffer from the ghost problem.

arXiv:1612.06104v1 [astro-ph.HE] 19 Dec 2016

The scenario — It was remarked that the observed orbital decay in a double neutron star (DNS) system, PSR 1913+16, is more rapid than that predicted by general relativistic quadrupole formula [1, 2], which was relieved by the scalar-tensor theory [3]. By making an analogy with the spontaneous magnetization of ferromagnets below the Curie temperature, an isolated neutron star (NS), with a compactness of $\frac{Gm}{Rc^2}$[1] above a certain critical value, will exhibit a nontrivial configuration, and a scalar field settles inside the NS [3, 4]. That is to say a spontaneous scalarization occurs for the NS. As a result, an external scalar field φ_1 appears around the scalarized NS, which triggers a scalarization of its companion, and subsequently another external scalar field φ_2 around the induced scalarized NS. The dynamical interplay between φ_1 and φ_2 is governed by a relation [5]

$$^{(n+1)}\varphi^i_B =^{(0)} \varphi^i + \frac{^{(n)}\varphi^j}{r}, \tag{1}$$

where the indices $i, j = 1, 2$ denote two NSs, and φ^i_B is the background field into which the NS i dips, $^{(0)}\varphi^i$ is the initial external scalar field produced by the NS i, $^{(n)}\varphi^j$ represents the nth induced scalar field around the NS j, and r is the distance from the centre of the binary. The feedback mechanism described by Eq. (1) results in an iteratively induced scalarization between two NSs, which enhances the strength of the external scalar fields. Accordingly, a convergence of φ^1_B and φ^2_B occurs, leading to a gravitational scalar background field ϕ. Therefore, the DNS is immersed in the gravitational scalar field ϕ, and the dynamics deviates from Einstein's general relativity, which imposes an influence on the orbital evolution of the binary system [6–8].

The dynamics of scalarized DNS binary is encoded not only by the gravitational tensor metric, but also by a gravitational scalar field. Scalar-tensor theory [6] of gravity naturally becomes the alternative theory to Einstein's general relativity describing the scalarized binary system. In this paper, we consider a minimally coupled scalar-tensor theory and investigate the mechanism that the spin-2 gravitons acquire a mass term due to the mediation of gravitational scalar field in scalarized DNS binaries. Instead of the usual method of effective field theory [9] that the graviton becomes heavy by eating Goldstone bosons because of the break of general coordinate invariance, we consider a potential, arising from the dynamical coupling between the gravitational scalar background field ϕ and the external scalar fields φ^i_B of each NS, as well as the self-coupling of ϕ. The iterative interplay between φ^1_B and φ^2_B causes small scalar fluctuations σ ($\sigma \ll \phi$), which in turn imposes a conformal transformation on both the tensor metric and the gravitational scalar field ϕ. By applying the conformal transformation to the tensor metric and scalar field that mediate the theory, the minimal coupling remains conformal invariant. The external background field φ^i_B with mass dimension, under the conformal transformation, introduces a mass dimensional constant, which is responsible for a spontaneous symmetry breaking. As a consequent, the gravitational scalar field obtains a mass, in which the fluctuation field σ plays the role of the Higgs field. The massive scalar field contributes to Yukawa type of corrections to the Newtonian potential of DNS binary, which, on the one hand, imposes a suppression on the propagation of massive scalar mode, and, on the other hand, influences the propagation of gravitational waves, endowing the gravitons with a mass.

The action — Neglecting the matter fields outside the NSs, we consider the scalar-tensor action that describes a DNS binary in vacuum,

$$S = \int d^4x\sqrt{-g}(\frac{M^2_{pl}}{2}\mathcal{R} - \frac{1}{2}g^{\mu\nu}\partial_\mu\phi\partial_\nu\phi - V(\phi)). \tag{2}$$

*Email address: wangjing6@mail.sysu.edu.cn

[1] m and R are the mass and radius of NS, respectively. G is the Newtonian gravitational constant.

Here, $M_{pl} = \sqrt{1/8\pi G}$ is the reduced Planck constant. $\mathcal{R}$ and g are the Ricci scalar and the determinant of the gravitational tensor metric $g_{\mu\nu}$, respevtively. $V(\phi)$ is the potential, consisting of the dynamical coupling of ϕ to φ_B^i and a self-coupling term of ϕ,

$$V(\phi) = \frac{\alpha}{2}(\varphi_B^i)^2\phi^2 + \frac{\lambda}{4}\phi^4. \tag{3}$$

Here, α is a dimensionless coupling constant and characterizes the coupling strength between the scalarized NS and ϕ, whose value depends on the compactness of the NSs [3, 4, 6]. λ is the self-coupling constant, which is roughly of the order of unity.

Considering the coupling between the scalar fluctuation σ and the dynamical fields, i.e. the tensor metric and the gravitational scalar field, we introduce a transformation [10]

$$\Omega(\sigma) = e^{\lambda\sigma} \quad (\sigma \ll 1), \tag{4}$$

The metric is transformed according to

$$g^*_{\mu\nu} = \Omega(\sigma)^{-2} g_{\mu\nu}, \quad \sqrt{-g^*} = \Omega(\sigma)^4\sqrt{-g}. \tag{5}$$

By expanding the new metric $g^*_{\mu\nu}$ about a Minkowski background in terms of Eq. (4) and Eq. (5), we express them as

$$g^*_{\mu\nu} = \eta_{\mu\nu} + h^*_{\mu\nu}, \quad h^*_{\mu\nu} = h_{\mu\nu} + 2\eta_{\mu\nu}\lambda\sigma, \tag{6}$$

where $|h_{\mu\nu}|, |h^*_{\mu\nu}| \ll 1$. The Eq. (5) remains unchanged. Applying Eq. (4) to ϕ, we get a new scalar field

$$\phi^* = \Omega(\sigma)^{-1}\phi, \tag{7}$$

and the kinetic term in action (2) is transformed into a canonical kinetic term,

$$-\frac{1}{2}\sqrt{-g}g_{\mu\nu}\partial_\mu\phi\partial_\nu\phi = -\frac{1}{2}\sqrt{-g^*}g^*_{\mu\nu}\mathcal{D}_\mu\phi^*\mathcal{D}_\nu\phi^*, \tag{8}$$

$$\mathcal{D}_\mu \equiv \partial_\mu + \lambda\partial_\nu\sigma, \tag{9}$$

which is scale-invariant. The transformed action then reads

$$S^* = \int d^4x\sqrt{-g^*}(\frac{M_{pl}^2}{2}\mathcal{R}^* - \frac{1}{2}g^{*\mu\nu}\mathcal{D}_\mu\phi^*\mathcal{D}_\nu\phi^* - V(\phi^*)). \tag{10}$$

The mass of gravitational scalar field — The solutions of external scalar fields φ_B^i for each component have mass dimensions [11]. In the process of conformal transformation, the solution with mass dimension involves a dimensional constant μ, which appears in the transformed scalar potential $V(\phi^*)$,

$$V(\phi^*) = \frac{\alpha}{2}\mu^2\phi^{*2} + \frac{\lambda}{4}\phi^{*4}. \tag{11}$$

The Planck scale constant $\mu = \sqrt{1/8\pi G_{\mathit{eff}}}$ appears to be related to the scalar charges of two scalarized NSs by the effective gravitational constant G_{eff} [6]. It is the appearance of the mass-dimensional constant μ that contributes to a process of spontaneous breaking of symmetry, which allows us to apply the similar recipe to the Higgs mechanism in the standard model.

Actual NSs observed in DNS binaries, with important deviations from general relativity in strong-field regime, would develop strong scalar charges in the absence of an external scalar field for enough negative values of α, i.e. $\alpha < 0$ [3–5]. The self-coupling constant λ is of the order of unity, i.e. $\lambda > 0$. By considering that the interplay between φ_B^i is long-range force, the behavior of the gravitational scalar field ϕ^* near spatial infinity endows it with a vacuum expectation value (VEV)

$$\phi_0^2 = -\frac{\alpha\mu^2}{2\lambda}, \tag{12}$$

which is obtained from the condition $\frac{dV(\phi^*)}{d\phi^*}|_{\phi^*_{min}} = 0$. Expanding around the VEV (12), we write the gravitational scalar background field as

$$\phi^* = \phi_0 + \tilde{\phi}. \tag{13}$$

Substituting the VEV (12) and Eq. (13) into the Lagrangian of ϕ^* extracted from Eq. (10), we can get the mass of ϕ^*,

$$m_s^2 = -\alpha\mu^2. \tag{14}$$

The scalar-mediated mass term of gravitons — Variation of the transformed action (10), with respect to the metric (5) and the new scalar field (7), yields the following equations of motion (e.o.m.) in vacuum,

$$\begin{aligned}(\mathcal{R}^*_{\mu\nu} - \frac{1}{2}g^*_{\mu\nu}\mathcal{R}^*)M_{pl}^2 &= \partial_\mu\phi^*\partial_\nu\phi^* - \frac{1}{2}g^*_{\mu\nu}(\partial_\alpha\phi^*)^2 \\ &\quad + \frac{\alpha}{2}\mu^2 g^*_{\mu\nu}\phi^{*2} + \frac{\lambda}{4}g^*_{\mu\nu}\phi^{*4} \quad (15)\\ \Box_{g^*}\phi^* &= \alpha\mu^2\phi^* + \lambda\phi^{*3}, \quad (16)\end{aligned}$$

where $\Box_{g^*}$ is the curved space d'Alembertian that is defined by $\Box_{g^*} = \sqrt{-g^*}\partial_\nu(\sqrt{-g^*}g^*_{\mu\nu}\partial_\mu)$.

Let us expand the transformed scalar potential $V(\phi^*)$ of Eq. (11) in a Taylor series about the VEV of ϕ^* (12),

$$V(\phi^*) = V(\phi_0) + V(\phi_0)'\tilde{\phi} + \frac{1}{2}V(\phi_0)''\tilde{\phi} + \cdots. \tag{17}$$

Considering the weak field perturbations of $g^*_{\mu\nu}$ expressed by Eq. (6) and the gravitational scalar field Eq. (13), we expand the field equations in weak field limit,

$$\begin{aligned}(\mathcal{R}^*_{\mu\nu} - \frac{1}{2}g^*_{\mu\nu}\mathcal{R}^*)M_{pl}^2 &= \partial_\mu\phi^*\partial_\nu\phi^* - \frac{1}{2}g^*_{\mu\nu}(\partial_\alpha\phi^*)^2 \\ &\quad + \frac{1}{2}m_s^2(\phi^* - \phi_0)^2 g^*_{\mu\nu}, \quad (18)\\ (\Box_g - m_s^2)\phi^* &= m_s^2\phi_0. \quad (19)\end{aligned}$$

Here, we use the expansion Eq. (17) and just consider the leading-order terms. The expanded field equations that are consistent at all orders in $(\frac{v}{c})^n$ is required. We solve the e.o.m. of massive scalar field Eq. (19) in a static spherically symmetric configuration, which yields an exterior solution of ϕ^* far from the DNS system,

$$\phi^* = \phi_0 + \phi_0 G_{\mathit{eff}} \frac{M_r}{r} e^{-m_s r}, \tag{20}$$

where $M_r = \frac{m_1 m_2}{m_1+m_2}$ is the reduced mass of DNS, and m_1 and m_2 are the masses of two NSs in the binary. Therefore, the mass m_s is the ordinary mass parameter in the Klein-Gordon equation (19) of ϕ^*, arising from the Higgs-like potential (11). In the meanwhile, the mass of ϕ^* plays precisely a role in the Yukawa-like correction $\sim e^{-m_s r}$ to the standard Newtonian form of gravitational potential $\sim \frac{GM_r}{r}$ in the scalarized DNS binary system.

In order to obtain the e.o.m. of gravitons, we expand the left-hand side of Eq. (18) in weak field limit, by using the weak field perturbations of tensor metric Eq. (6). Imposing the harmonic gauge $\partial^\nu(h^*_{\mu\nu} - \frac{1}{2}\eta_{\mu\nu}h^*) = 0$, with $h^* = \eta^{\mu\nu}h^*_{\mu\nu}$, and neglecting the higher-order terms, we rewrite the e.o.m. of gravitons as

$$\begin{aligned} -\frac{M_{pl}^2}{2}\Box_\eta \bar{h}^*_{\mu\nu} \;=\; & \partial_\mu\phi^*\partial_\nu\phi^* - \frac{1}{2}\eta_{\mu\nu}(\partial_\alpha\phi^*)^2 \\ & + \frac{1}{2}m_s^2(\phi^* - \phi_0)^2\eta_{\mu\nu}, \end{aligned} \tag{21}$$

where $\bar{h}^*_{\mu\nu} = h^*_{\mu\nu} - \frac{1}{2}\eta_{\mu\nu}h^*$ and the flat-space d'Alembertian $\Box_\eta = \eta^{\mu\nu}\partial_\mu\partial_\nu$. Let us study the scalar-mediated propagations of gravitons outside the scalarized DNS binary. We substitute the solution of ϕ^* (20) into the e.o.m. of gravitons (21) and follow the gauge selection described in [12]. The e.o.m. of gravitons (21) has the wave solutions

$$\bar{h}^*_{\mu\nu} = \Big(\int d\omega \int \frac{d^3k}{(2\pi)^3} A e^{i(\vec{k}\cdot\vec{r}-\omega t)}\Big) \cdot \sum_{n=-\infty}^{\infty} \phi^{*n}(\vec{r}) e^{in\sigma}. \tag{22}$$

Here, A denotes the amplitude of tensor gravitational waves radiated from the orbital decaying DNS. $\sum_{n=-\infty}^{\infty} \phi^{*n}(\vec{r})e^{in\sigma}$ is the Fourier expansion of the gravitational scalar field ϕ^*, with the fluctuation fields σ, which is converged by the nth induced scalar background ${}^n\varphi^i_B$ of two NSs. ϕ^{*n} is in the form of exterior solutions (20). From Eq. (22), we find that the propagation of graviton depends on five components, i.e. the space and time coordinates ($\vec{r}$ and t) in 4 dimensions, and a scalar component as the 5th dimension. The Klein-Gordon equation of gravitons (22) therefore reads

$$[\Box - (k^2 - \omega^2) - \frac{m_s^2\phi_0 G_{\mathit{eff}}}{\tilde{R}}]\bar{h}^*_{\mu\nu} = 0, \tag{23}$$

where $\tilde{R}$ is the semi-major axis of the elliptical DNS binary system. By defining

$$m_g^2 = \frac{m_s^2\phi_0 G_{\mathit{eff}}}{\tilde{R}}, \tag{24}$$

we find that the gravitons acquire a mass of m_g. In the scenario, the massive gravitational scalar field ϕ^* modifies the Newtonian potential of DNS and contributes to a Yukawa-like one $\sim \frac{e^{-m_s r}}{r}$, which imposes influence on the propagations of tensor gravitational waves, via the entrance of massive scalar component into the propagation of gravitational waves. Hence, the gravitons acquire a mass. The scalar fluctuation σ is the only massless field, which plays the role of Higgs-like field.

Ghost-free arguments — The gravitons, with mass degree of freedom (d.o.f.), generally suffer from a central problem – ghost instability, which has been widely discussed and attempted to solve in many ways [13]. When we study the wave solutions for the propagation of $h^*_{\mu\nu}$, the four constraint equations [12] along with four general coordinate transformations generically remove the four non-propagating components of the metric, with the broken general covariance, remaining six propagating modes. However, the massive gravitons have five d.o.f., corresponding to five physical modes of polarizations. The remaining mode is a ghost. The e.o.m. of gravitons (21), under the transverse-traceless gauge condition, propagates six d.o.f., which contains a massive scalar mode. From the wave solutions of gravitons (22), the gravitational waves propagate in a 5-dimension-like space-time near the binary system ($r \sim 10^9$ m). The appearance of massive gravitational scalar field modifies the interaction between two NSs and produces a Yukawa-like potential $\sim e^{m_s r}/r$, which makes the massive scalar gravitational field ϕ^* exponentially decrease outside the DNS binary. That is to say the significant effects of the massive scalar field and the propagation of massive mode is screened in a scale of binary system. The screening scale $\tilde{R}$ in turn gives the massive scalar mode a characteristic range $\tilde{R}^{-1}$. The effects of Yukawa-like potential enter the solutions of gravitational waves via the massive gravitational scalar field and thus influence the propagations of $h^*_{\mu\nu}$ in the near region of the binary, which endows the gravitons with a mass. On the scale of DNS binary ($r \sim 10^9$ m), the radiated gravitational waves propagate six modes, i.e. massive graviton with five physical d.o.f. and a massive scalar mode that is screened in the binary system. While on a large scale outside the binary ($r \gg 10^9$ m), the gravitational waves propagate with five modes of polarizations, corresponding to ghost-free massive gravitons.

Constraints from detected DNS binaries — According to the expressions of (12) and (14), we rewrite the mass of gravitons, Eq.(24), as

$$m_g^2 = (-\alpha)^{3/2}\mu^3(2\lambda)^{-1/2}G_{\mathit{eff}}\tilde{R}^{-1}. \tag{25}$$

It is found that the mass of graviton depends on three quantities[2], i.e. the separation of DNS system represented by $\tilde{R}$, the coupling strength between gravitational scalar field and the NSs that is characterized by the dimensionless coupling constant α, and the scalar charges that is related to both the Planck scale constant μ and the effective gravitational constant G_{eff}. Consequently, the mildly varying mass of graviton rests with the intrinsic properties of the DNS, as well as the coupling strength between the NS matter and the gravitational scalar field.

It was proven that nonperturbative strong-gravitational-field effects developed in NSs for a dimensionless coupling constant $\alpha \lesssim -4$, which induced order-of-unity deviations from general relativity [3]. The general properties of binary systems consisting of scalarized NSs can be described by $\alpha \gtrsim -4.5$, because of binary-pulsar measurements [14–16]. For $\alpha \lesssim -5$, NSs in binary pulsar, with mass of 1.4 $M_\odot$, would develop strong scalar charges even in absence of external scalar solicitation, and a more negative value of α corresponds to a less compact NS [4]. Most of the measured more massive NS in detected DNS systems have masses of $\sim 1.3 - 1.44 M_\odot$ [17]. Consequently, the coupling constant locates in a range of $\alpha = -5 \sim -6$ within a quadratic coupling model described in Eq. (11) [4]. The scalar charges mildly vary with the compactness of NSs [5] and will be ~ 1 only in the last stages of the evolution of NS binaries or close transient encounters. For NSs in the 9 detected DNS systems, the scalar charges are around 0.2 within solar-system bound [18] in the Fierz-Jordan-Brans-Dicke (FJBD) theory, by considering its dependence on the "sensitivities" $s \sim 0.2$ [19, 20]. Accordingly, the gravitons radiated in a DNS binary with a semi-separation of 10^9 m have masses of $\sim 10^{-24} ev/c^2$, which mildly vary with the compactness of NSs and the separation between them.

i) PSR J1913+16 — The Hulse-Taylor system consists of two pulsars with masses of $m_1 = 1.44 \pm 0.0006 M_\odot$ and $m_2 = 1.39 \pm 0.0006 M_\odot$. The orbital period is 0.323 day, and an elliptical orbit has orbital eccentricity of 0.617. According to Eq. (25), we can get that the radiated gravitons have the mass of $m_g = 9.715 \times 10^{-24} ev/c^2$.

ii) PSR J0737-3039A(B) — Another DNS binary system, consisting of two pulsars, has a closed and approximate circular orbit with an eccentricity of 0.088. The orbital period is 0.102 day. The measured masses of two radio pulsars are $m_1 = 1.34 \pm 0.010 M_\odot$ and $m_2 = 1.25 \pm 0.010 M_\odot$, respectively. Accordingly, we get the radiated gravitons have mass of $3.356 \times 10^{-23} ev/c^2$.

iii) PSR B1534+12 — An approximate equal-mass DNS binaries with NS masses of $m_1 = 1.35 \pm 0.0020 M_\odot$ and $m_2 = 1.33 \pm 0.0020 M_\odot$. The two NSs orbit with each other at an orbital period of 0.421 day on an elliptical orbit with an eccentricity of 0.274. The mass of graviton thus is $1.292 \times 10^{-23} ev/c^2$.

iv) PSR B2127+11C — Another approximate equal-mass DNS binaries with NS masses of $m_1 = 1.35 \pm 0.080 M_\odot$ and $m_2 = 1.36 \pm 0.080 M_\odot$. The two NSs orbit with each other at an orbital period of 0.335 day on an eccentric orbit with eccentricity of 0.681. The estimated mass of graviton is $1.408 \times 10^{-23} ev/c^2$.

v) PSR J1829+2456 — A DNS with an orbital period of 1.176 day shows low eccentricity of 0.139. The masses of two NSs are $m_1 = 1.15^{+0.1}_{-0.25} M_\odot$ and $m_2 = 1.35^{+0.46}_{-0.15} M_\odot$, respectively. Within the quadratic coupling potential described by Eq. (11), the coupling constant is $\alpha \sim -6$. Consequently, the mass of gravitons is estimated as $7.106 \times 10^{-24} ev/c^2$.

vi) PSR J1906+0746 — An approximate circular DNS system have the similar properties with the system PSR J0737-3039, which shows orbital period of 0.166 day. Accordingly, the mass of gravitons is $2.427 \times 10^{-23} ev/c^2$.

vii) PSR J1811-1736 — A very wide DNS binary has an orbital period of 18.8 days and a very eccentric orbit with an eccentricity of 0.828. The two NSs have masses of $m_1 = 1.5^{+0.12}_{-0.4} M_\odot$ and $m_2 = 1.06^{+0.45}_{-0.1} M_\odot$. The gravitons from the system is $m_g = 1.063 \times 10^{-24} ev/c^2$.

viii) PSR J1756-2251 — The system has NSs masses of $m_1 = 1.40^{+0.04}_{-0.06}$ and $m_2 = 1.18^{+0.06}_{-0.04}$, an orbital period of 0.320 days, and an eccentricity of 0.181. The mass of gravitons radiated from this system is $1.574 \times 10^{-23} ev/c^2$.

Final remarks — When we estimate the mass of gravitons for the above 8 detected DNS binaries, we use the assumption of a quadratic coupling between the scalar field and NS matter. If the coupling form is changed, we may get a different coupling strength, as well as the mass of gravitons. Therefore, the value of graviton mass in 8 DNS systems we estimated is model-dependent.

This work is supported by the Fundamental Research Funds for the Central Universities (Grant no. 161gpy49) at Sun Yat- Sen University.

[2] Noting that the self-coupling constant λ of gravitational scalar field is of the order of unity.

[1] J. H. Taylor and J. M. Weisberg, "A new test of general relativity - Gravitational radiation and the binary pulsar PSR 1913+16," Astrophys. J., 253, 908 (1982)

[2] J. M. Weisberg, D. J. Nice and J. H. Taylor, "Timing Measurements of the Relativistic Binary Pulsar PSR B1913+16," Astrophys. J., 722, 1030 (2010)

[3] T. Damour and G. Esposito-Farese, "Nonperturbative strong field effects in tensor - scalar theories of gravitation," Phys. Rev. Lett. **70**, 2220 (1993).

[4] T. Damour and G. Esposito-Farese, "Tensor - scalar gravity and binary pulsar experiments," Phys. Rev. D **54**, 1474 (1996)

[5] C. Palenzuela, E. Barausse, M. Ponce and L. Lehner, "Dynamical scalarization of neutron stars in scalar-tensor gravity theories," Phys. Rev. D **89**, no. 4, 044024 (2014)

[6] T. Damour and G. Esposito-Farese, "Tensor multiscalar theories of gravitation," Class. Quant. Grav. **9**, 2093 (1992).

[7] R. A. Hulse and J. H. Taylor, "Discovery of a pulsar in a binary system," Astrophys. J. Lett., 195, L51 (1975)

[8] C. M. Will and H. W. Zaglauer, "Gravitational radiation, close binary systems, and the Brans-Dicke theory of gravity," Astrophys. J, 346, 366 (1989)

[9] N. Arkani-Hamed, H. Georgi and M. D. Schwartz, Annals of Physics, 305, 96 (2003)

[10] Y. Fujii and K. Maeda, "The scalar-tensor theory of gravitation" Cambridge University Press, Cambridge, England, (2007)

[11] M. Salgado, D. Sudarsky and U. Nucamendi, "On spontaneous scalarization," Phys. Rev. D **58**, 124003 (1998) doi:10.1103/PhysRevD.58.124003 [gr-qc/9806070].

[12] Michele Maggiore, "Gravitational Waves Volume 1: Thoery and Experiments" Oxford University Press, London, England, (2008)

[13] C. de Rham, "Massive Gravity," Living Reviews in Relativity, 17, 7 (2014)

[14] T. Damour and G. Esposito-Farese, "Gravitational wave versus binary - pulsar tests of strong field gravity," Phys. Rev. D **58**, 042001 (1998) [gr-qc/9803031].

[15] P. C. C. Freire, N. Wex, G. Esposito-Farèse, J. P. W. Verbiest, M. Bailes, B. A. Jacoby, M. Kramer, I. H. Stairs, J. Antoniadis, and G. H. Janssen, "The relativistic pulsar-white dwarf binary PSR J1738+0333 - II. The most stringent test of scalar-tensor gravity," Mon. Not. R. Astron. Soc., 423, 3328 (2012)

[16] J. Antoniadis, P. C. C. Freire, N. Wex, T. M. Tauris, R. S.Lynch, M. H. van Kerkwijk,, M. Kramer, C. Bassa, V. S. Dhillon, T. Driebe, J. W. T. Hessels, V. M. Kaspi, V. I. Kondratiev, N. Langer, T. R. Marsh, M. A. McLaughlin, T. T. Pennucci, S. M. Ransom, I. H. Stairs, J. van Leeuwen, J. P. W. Verbiest, D. G. Whelan, "A Massive Pulsar in a Compact Relativistic Binary," Science, 340, 448 (2013)

[17] C. M. Zhang, J. Wang, Y.-H. Zhao, H. X. Yin, L. M. Song, D. P. Menezes, D. T. Wickramasinghe, L. Ferrario, P. Chardonnet, "Study of measured pulsar masses and their possible conclusions," Astron. Astrophys., 527, A83 (2011)

[18] D. M. Eardley, "Observable effects of a scalar gravitational field in a binary pulsar," Astrophys. J. Lett., 196, L59 (1975)

[19] T. Damour and G. Esposito-Farese, "Testing gravity to second postNewtonian order: A Field theory approach," Phys. Rev. D **53**, 5541 (1996) [gr-qc/9506063].

[20] S. Mirshekari and C. M. Will, "Compact binary systems in scalar-tensor gravity: Equations of motion to 2.5 post-Newtonian order," Phys. Rev. D **87**, no. 8, 084070 (2013) [arXiv:1301.4680 [gr-qc]].